Front Line Demonstration (Conducted during 2024)

Title	Technology	Photo
Popularization of blackgram var. PU-31 with improved agronomic practices	 Popularization of blackgram variety PU-31 Seed rate: 20 kg/ha Spacing: 30cm x10 cm Seed treatment with Carbendazim+Mancozeb @ 2g/kg seed NPKS@20:40:20:15 kg/ha Seed inoculation with Rhizobium@50g+10 g sugar per kg seed 	Utlou, Manipur, India Pvf5+g53, Utlou, Yarou Meitram, Manipur 795134, India Lat 24.723841° Long 93.859159° 08/11/24 12:32 PM GMT +05:30
Popularization of Broccoli cultivation intercropped with Coriander	 ▶ Broccoli var. Green magic intercropped with coriander ▶ Seed rate of Broccoli - 350g/ha, ▶ Spacing-60cmx45 cm, ▶ Seed rate of coriander 10 kg/ha ▶ Application of Vermicompost- 5 tons/ha 	Though Like Branch Paris Andrews Haward Like Branch Paris Andrews Like Branch Paris Andrews Like Branch Paris Haward Like Branch Paris Andrews Like Branch Paris Haward Like Branch Paris Like B
Popularization on Scientific cultivation of Onion var. Arka Kalyan	 ➢ Onion : Var. Arka Kalyan ➢ Seed rate : 10 kg/ha, ➢ Spacing : 15 cmx10 cm ➢ FYM : 5 tons/ha ➢ NPK : 120:60:40kg/ha 	Toubul, Manipur, भारत Toubul Road, Toubul, Manipur 795126, भारत Lat 24.624799° Long 93.78362° 08/04/2025 11:56 AM GMT +05:30
Production technology of Stunted fish fingerlings.	Stocking of IMC Fish fry at high stocking density @2,50,000 fry /ha for 10 months with natural feed	
Popularisation of jackfruit chip	Preparation with blanching ❖ Cutting of fully matured unripe jackfruit. Peeling and deseeding ❖ Deseeded bulbs cutting longitudinal into finger like pieces ❖ Blanching into hot water with 1% KMS for 5 minutes ❖ Dried in drier @ 42°C	
Management of fall armyworm in maize.	Application of Emamectin benzoate 5SG @0.4g/l at the interval of 10 days.	KRISHIVGAN KEADRA Front Line Demonstration on —Management of Fall Armyworm in Male Lecture Start
Integrated pest and disease management in pea.	Pre sowing- summer plough, removal of residue of previous crop. Seedling & veg stage: collection and destruction of insects larvae. Seed treatment with Trichoderma viridae @5-10/kg before sowing.& application of neem oil 0.15 EC @ 3ml/l to control aphids.	Front Line Demonstration on Integrated Pest & Disease Management in Pea Location: Bishnupur Year: 2024
Popularization of Low cost Vermicomposting technology		
Popularization of Enriched compost (Made from locally available biomass)		REISHI VIGYAN KENDRA Bilitarque Detrit, Manjuer Front Line Demonstration on Cornect vom hearly variable tolerance Acre Dos 100 Dos 10